

# On Boffa Alternative

B. Bajorska, O. Macedońska

August 17, 2013

## Abstract

Let  $G^*$  denote a nonprincipal ultrapower of a group  $G$ . In 1986 M. Boffa posed a question equivalent to the following one: *if  $G$  does not satisfy a positive law, does  $G^*$  contain a free nonabelian subsemigroup?* We give the affirmative answer to this question in the large class of groups, containing all residually finite and all soluble groups, in fact all groups, considered in traditional textbooks on group theory.

**Preliminaries** Let  $\mathcal{F}$  be a free semigroup generated by  $x_1, x_2, \dots$ . We say that an  $n$ -tuple of elements  $g_1, \dots, g_n$  in a group  $G$  satisfies a nontrivial positive relation  $u(x_1, \dots, x_n) = v(x_1, \dots, x_n)$ , if  $u, v$  are different words in  $\mathcal{F}$  and the equality  $u(g_1, \dots, g_n) = v(g_1, \dots, g_n)$  holds. A group  $G$  satisfies the law  $u = v$  if the equality holds under each substitution of elements in  $G$  for generators  $x_i$ . We note that every  $n$ -variable positive law implies a nontrivial two-variable positive law if we replace each variable  $x_i$  by  $xy^i$ .

## Notation

We say that  $G$  is an  $\mathcal{F}$ -group (respectively an  $\mathcal{F}^*$ -group) if  $G$  (resp.  $G^*$ ) contains a free nonabelian subsemigroup;

We say that  $G$  is a  $no\mathcal{F}$ -group (resp. a  $no\mathcal{F}^*$ -group) if  $G$  (resp.  $G^*$ ) does not contain a free nonabelian subsemigroup.

Since  $G^*$  is the image of a cartesian power  $G^\omega$  under the congruence defined by a nonprincipal ultrafilter, the group  $G$  is embedded into  $G^*$  via  $diag G^\omega$ . Hence every  $\mathcal{F}$ -group is an  $\mathcal{F}^*$ -group. To see that the class of  $\mathcal{F}^*$ -groups is wider, we note that in a  $no\mathcal{F}^*$ -group  $G$  every pair of elements satisfies a two-variable relation from some finite set  $\{u_i = v_i, i = 1, 2, \dots, n\}$  (see Lemma

---

2000 AMS Subject Classification 20F19, 20E10

2 below), and hence  $G$  satisfies a nontrivial (left-normed) commutator law  $[u_1v_1^{-1}, u_2v_2^{-1}, \dots, u_nv_n^{-1}]$ . So infinite residually finite torsion groups which by [12], IV.2.6, do not satisfy any law, are  $\mathcal{F}^*$ -groups.

Question (Q) [1] suggests the following

**Boffa alternative: A group either satisfies a positive law or is an  $\mathcal{F}^*$ -group.**

Since nilpotent groups satisfy a positive law [7], every nilpotent-by-(finite exponent) also does, hence trivially satisfies the Boffa alternative. So finitely generated soluble groups ([8], Thms. 4.7, 4.12), elementary amenable groups [3], finitely generated linear groups over any field [10] satisfy the Boffa alternative. Every relatively free group satisfies Boffa alternative because every positive relation on free generators provides a positive law.

It is not known whether every group satisfies the alternative. In this paper we present a large class of groups satisfying the Boffa alternative. We need the following

**Notation** Let  $\mathfrak{N}_c$  denote the variety of all nilpotent groups of nilpotency class  $c$ ,  $\mathfrak{S}_n$  the variety of all soluble groups of solubility class  $n$ , and  $\mathfrak{B}_k$  the restricted Burnside variety of exponent  $k$ , that is, the variety generated by all finite groups of exponent  $k$ . It follows from the positive solution of the Restricted Burnside Problem (see [11]) that all groups in  $\mathfrak{B}_k$  are locally finite of exponent dividing  $k$ .

To recall the definition of the class  $\mathcal{C}$  introduced in [2], we denote by  $\Delta_1$  the class of groups contained in all finite products  $\mathfrak{V}_1\mathfrak{V}_2\dots\mathfrak{V}_m$ , where  $\mathfrak{V}_i$  is either  $\mathfrak{S}_n$  or  $\mathfrak{B}_k$  for various  $n, k$ . Now we define inductively:  $\Delta_{n+1} = \{\text{groups, locally in } \Delta_n\} \cup \{\text{groups, residually in } \Delta_n\}$ .

The class  $\mathcal{C}$  is defined as the union:  $\mathcal{C} = \cup_n \Delta_n$ .

Among all laws satisfied in  $G \in \mathfrak{N}_c\mathfrak{B}_k$  there is a two-variable positive law of the form  $u_c(x, y) = u_c(y, x)$ , where  $u_1(x, y) = x^k y^k$ , and  $u_c(x, y) = u_{c-1}(x, y)u_{c-1}(y, x)$  (cf. [7]). For groups with certain properties the parameters  $c, k$  can be bounded as functions of some  $n$ . Then we denote such a law by  $M_{ck}(n)$ . We need the following

**Lemma 1** (cf. [9], p.52, Thms. A', B) *Let  $G$  be a two-generator residually finite group, and let there exist  $n$  such that for every  $n$ -element subset  $S \subseteq G$ , the inequality  $|S^n| < n^n$  holds. Then  $G$  satisfies the law  $M_{ck}(n)$ , where  $c, k$  depend on  $n$  only.  $\square$*

### Properties of $no\mathcal{F}^*$ -groups

By [1], Thm. 3,  $G$  is a  $no\mathcal{F}^*$ -group if and only if then there exists a finite set  $\mathfrak{A}$  of nontrivial two-variable positive relations such that every pair of elements in  $G$  satisfies at least one of them.

We denote by  $\mathfrak{L}_n$  the set of all nontrivial two-variable positive relations of the length  $n$ , that is:

$$\mathfrak{L}_n = \{u(x, y) = v(x, y); |u(x, y)| = |v(x, y)| = n\},$$

where  $|u(x, y)|$  is the length of the word  $u(x, y)$ .

The following is clear: a relation  $u = v$  implies both  $uv = vu$  and  $xu = xv$ ; a relation  $uvw = w$  implies  $uv = vu$ . Now it can be easily shown that for any finite set  $\mathfrak{A}$  of nontrivial two-variable positive relations there exists  $n$  such that: if every pair of elements in  $G$  satisfies at least one relation in  $\mathfrak{A}$ , then every pair of elements in  $G$  satisfies at least one relation in  $\mathfrak{L}_n$ . So we obtain

**Lemma 2** (cf. [1], Thm. 3) *A group  $G$  is a  $no\mathcal{F}^*$ -group if and only if there exists  $n$  such that every pair of elements in  $G$  satisfies at least one relation in  $\mathfrak{L}_n$ .  $\square$*

From the above Lemma it follows that for  $N \triangleleft H \subseteq G$ , every pair of elements in  $H/N$  also satisfies at least one relation in  $\mathfrak{L}_n$  and hence  $H/N$  is a  $no\mathcal{F}^*$ -group.

Now we need to show that if every pair of elements in  $G$  satisfies at least one relation in  $\mathfrak{L}_n$ , then every residually finite subgroup in  $G$  and every residually finite quotient of  $G$  satisfy the positive law  $M_{ck}(n)$ , where the parameters  $c, k$  depend on  $n$  only.

**Lemma 3** *Let  $G$  be a  $no\mathcal{F}^*$ -group and  $N \triangleleft H \subseteq G$ . If  $H/N$  is locally residually finite, then there exists  $n$  such that  $H/N$  satisfies the law  $M_{ck}(n)$ .*

**Proof** If  $h_1, h_2$  are elements in  $H/N$ , then by assumption, the subgroup  $gp(h_1, h_2)$  is residually finite. By Lemma 2, there exists  $n$  such that every pair of elements in  $G$ , and hence in  $H/N$ , satisfies a relation in  $\mathfrak{L}_n$ . We consider any set  $S = \{s_1, \dots, s_n\}$  in  $gp(h_1, h_2)$ . Elements  $s_1, s_2$  satisfy some relation  $u(x, y) = v(x, y)$  in  $\mathfrak{L}_n$ , so the words  $u(s_1, s_2)$  and  $v(s_1, s_2)$  are in  $S^n$ , and are equal. Thus we have  $|S^n| < n^n$ , and by Lemma 1, the group  $gp(h_1, h_2)$  satisfies the law  $M_{ck}(n)$ . So the whole section  $H/N$  satisfies the law  $M_{ck}(n)$ , as required.  $\square$

**Corollary 1** *If  $G$  is a  $no\mathcal{F}^*$ -group, then there exists  $n$  such that every soluble subgroup  $H$  in  $G$  satisfies the law  $M_{ck}(n)$ .*

**Proof** By Lemma 2, there exists  $n$  such that every pair of elements in  $G$  (and hence in  $H$ ) satisfies a relation in  $\mathfrak{L}_n$ . Let  $h_1, h_2$  be in  $H$ . Then the subgroup  $gp(h_1, h_2)$  is a finitely generated, soluble  $no\mathcal{F}$ -group. Now by Thms. 4.7 and 4.12 in [8],  $gp(h_1, h_2)$  is an extension of a nilpotent group by a finite group, and hence by [4], is residually finite. So by Lemma 3, the elements  $h_1, h_2$  satisfy  $M_{ck}(n)$ . Since the same is true for any  $h_1, h_2$  in  $H$ , we conclude that  $H$  satisfies the law  $M_{ck}(n)$ .  $\square$

### The Main Result

To show that every group in the class  $\mathcal{C}$  satisfies the Boffa alternative, it is enough to prove that every  $no\mathcal{F}^*$ -group in the class  $\mathcal{C}$  satisfies a positive law.

We start with a  $no\mathcal{F}^*$ -group in a product of a restricted Burnside variety and a soluble variety.

**Lemma 4** *If  $G$  is a  $no\mathcal{F}^*$ -group in  $\mathfrak{B}_e\mathfrak{S}_d$ , then  $G$  satisfies a positive law, and  $G \in \mathfrak{N}_c\mathfrak{B}_k$  for some  $c, k$*

**Proof** By Lemma 2, there exists  $n$  such that every pair of elements in  $G$  satisfies a relation in  $\mathfrak{L}_n$ . We show that  $G$  satisfies the law  $M_{ck}(n)$ . If  $H$  is a finitely generated subgroup in  $G$ , then  $H$  is a  $no\mathcal{F}$ -group and by ([6], Corollary 3), all its derived subgroups are finitely generated. Since by assumption  $H \in \mathfrak{B}_e\mathfrak{S}_d$ , we get  $H^{(d)} \in \mathfrak{B}_e$ . So  $H^{(d)}$  is a finitely generated group in  $\mathfrak{B}_e$  and hence is finite. The centralizer  $Z$  of the finite subgroup  $H^{(d)}$  in  $H$  must have a finite index in  $H$  ([5], 3.1.4), and hence is finitely generated ([5], 14.3.2). Moreover,  $Z$  is soluble, because  $1 = [H^{(d)}, Z] \supseteq [Z^{(d)}, Z^{(d)}] = Z^{(d+1)}$ .

The finitely generated soluble  $no\mathcal{F}$ -group  $Z$  is, by [8], nilpotent-by-finite. So  $H$ , as a finite extension of  $Z$ , is also nilpotent-by-finite. Now  $G$  locally is nilpotent-by-finite and hence by [4], locally residually finite. So by Lemma 3,  $G$  satisfies the positive law  $M_{ck}(n)$ . By Theorem B in [2] it follows that  $G$  is in  $\mathfrak{N}_c\mathfrak{B}_k$  for some  $c, k$ , as required.  $\square$

**Theorem 1** *If  $G$  is in the class  $\mathcal{C}$ , then either  $G$  satisfies a positive law or  $G$  is an  $\mathcal{F}^*$ -group.*

**Proof** It is enough to show that if  $G \in \mathcal{C}$  is a  $no\mathcal{F}^*$ -group then  $G$  satisfies a positive law. By Lemma 2 we assume that there exists  $n$  such that every pair of elements in  $G \in \mathcal{C}$  satisfies a relation in  $\mathfrak{L}_n$ . We shall prove that  $G$  satisfies the law  $M_{ck}(n)$ .

Let our group  $G$  be in  $\Delta_1$ . Then  $G \in \mathfrak{V}_1\mathfrak{V}_2 \dots \mathfrak{V}_m$ , where each variety  $\mathfrak{V}_i$  is either  $\mathfrak{S}_d$  or  $\mathfrak{B}_e$  for some  $d, e$ . Since the product of varieties is associative, we replace, by Lemma 4, (starting from the right) every pair of the type  $\mathfrak{B}\mathfrak{S}$  by some pair of the type  $\mathfrak{N}\mathfrak{B}$ , and find that  $G$  belongs to a soluble-by-restricted Burnside variety  $\mathfrak{S}_k\mathfrak{B}_e$  for some  $k, e$ . Then  $G$  is a locally soluble-by-finite  $no\mathcal{F}$ -group and by [8], locally is nilpotent-by-finite. Hence  $G$  locally is residually finite and then by Lemma 3, satisfies the law  $M_{ck}(n)$ .

So: if in  $G \in \Delta_1$  every pair of elements satisfies a relation in  $|\text{cal}L_n$ , then  $G$  satisfies the law  $M_{ck}(n)$ .

We now consider a group  $G$  in  $\Delta_2$ . Since  $\Delta_2 = \{\text{groups, locally in } \Delta_1\} \cup \{\text{groups, residually in } \Delta_1\}$ , there are two cases.

(i) If  $G$  is locally in  $\Delta_1$ , and if  $K$  is a finitely generated subgroup in  $G$ , then  $K \in \Delta_1$  and hence, as we have proved above, satisfies the law  $M_{ck}(n)$ . Since the same is true for every finitely generated subgroup  $K \subseteq G$ , we conclude that  $G$  satisfies the positive law  $M_{ck}(n)$ .

(ii) If  $G$  is residually in  $\Delta_1$ , then  $G$  is a subcartesian product of its finite quotients  $G/N \in \Delta_1$ . By assumption  $G$  and hence  $G/N$  are  $no\mathcal{F}^*$ -groups. Hence, as we have proved, each  $G/N$  satisfies the law  $M_{ck}(n)$ . So  $G$ , as a subcartesian product of  $G/N$ 's, satisfies the same law.

It follows that every  $no\mathcal{F}^*$ -group  $G$  in  $\Delta_2$  satisfies a positive law. Then by Theorem B in [2], there exist  $s, t$  such that  $G \in \mathfrak{N}_s\mathfrak{B}_t$ , and hence  $G$  is in  $\Delta_1$ , which ends the induction. So if  $G$  is an  $\mathcal{F}^*$ -group in the class  $\mathcal{C}$ , then  $G$  is in  $\Delta_1$  and as we have proved, satisfies a positive law.  $\square$

By Theorem B in [2] it follows that every group  $G \in \mathcal{C}$  either is in  $\mathfrak{N}_c\mathfrak{B}_k$  for some  $c, k$  or is an  $\mathcal{F}^*$ -group.

## References

- [1] M. Boffa, Elimination of inverses in groups, *Model Theory of Groups and Automorphism Groups, London Math. Soc. Lecture Notes Series*, **224** p. 134-143.

- [2] R. G. Burns, O. Macedońska, Yu. Medvedev, Groups satisfying semi-group laws, and nilpotent-by-Burnside varieties, *J. Algebra* **195** (1997), 510-525.
- [3] C. Chou, Elementary amenable groups, *Ill. J. Math.* **24** (1980), 396-407.
- [4] P. Hall, On the finiteness of certain soluble groups, *Proc. London Math. Soc.* **9** (1959), 595-622.
- [5] M. I. Kargapolov, Yu. I. Merzljakov, Fundamentals of the theory of groups, *Nauka, Moscow 1982*.
- [6] P. Longobardi, M. Maj, A. H. Rhemtulla, Groups with no free sub-semigroups, *Trans. Amer. Math. Soc.* **347** (1995), 1419-1427.
- [7] A. I. Mal'cev, Nilpotent semigroups, *Uchen. Zap. Ivanovsk. Ped. Inst.* **4** (1953), 107-111.
- [8] J. M. Rosenblatt, Invariant measures and growth conditions, *Trans. Amer. Math. Soc.* **193**, (1974), 33-53.
- [9] A. Shalev, Combinatorial conditions in residually finite groups, II, *J. Algebra* **157** (1993), 51-62.
- [10] J. Tits, Free subgroups in linear groups, *J. Algebra* **20** (1972), 250-270.
- [11] M. Vaughan-Lee, On Zelmanov's solution of the restricted Burnside problem, *J. Group Theory* **1** (1998), no. 1, 65-94.
- [12] E. I. Zelmanov, Nil Rings and Periodic Groups, *The Korean Mathematical Society, 1992*.

Institute of Mathematics,  
 Silesian University of Technology,  
 ul. Kaszubska 23,  
 44-100 Gliwice, Poland  
 e-mail: olga@zeus.polsl.gliwice.pl  
 e-mail: bajorska@zeus.polsl.gliwice.pl